

May the Force Be with Youth: Foundational Strength for Lifelong Development

Avery D. Faigenbaum, EdD, FACSM;¹ Nicholas A. Ratamess, PhD;¹ Jie Kang, PhD, FACSM;¹ Jill A. Bush, PhD, FACSM;¹ and Tamara Rial Rebullido, PhD²

Abstract

Today's youth are weaker than previous generations, and measurable reductions in physical fitness are beginning to emerge. Without targeted initiatives that recognize the foundational importance of resistance training, weaker children and adolescents may be more likely to experience the inevitable consequences of neuromuscular dysfunction and less likely to experience the pleiotropic benefits of exercise and sport. Early exposure to strength-building activities is needed to prepare today's youth for ongoing participation in varied physical activities throughout this developmental phase of life. The novel iceberg of physical development is a metaphoric image that illustrates the sequential and cumulative influence of muscular strength on motor skills and physical abilities. Efforts to enhance the physical capacity of youth should include resistive skills that improve basic movement patterns and enhance motoric competence. A shift in our conceptual thinking about youth resistance training is needed to alter the current trajectory toward physical inactivity and related comorbidities.

and force reduction (*e.g.*, landing from a jump) is needed to move and play skillfully, weaker youth may be less likely to accumulate at least 60 min of MVPA daily and more likely to experience functional limitations, activity-related injuries, and related comorbidities (10–12).

Measurable reductions in physical fitness are emerging, sport participation rates during adolescence are declining, and the medical and financial burden of activity-related injuries in young athletes are substantial (2,13,14). Developmentally appropriate interventions that recognize the foundational importance of muscular fitness are needed to prepare today's youth for the demands of rough and tumble play, vigorous exercise, and sports competition (11,15,16). Low levels of muscular strength during childhood tend to track into adulthood (17,18), and therefore weaker children may not simply “out-grow” strength deficits. An integrative approach that develops biomotor abilities including muscular strength, speed, and coordination is needed before youth become disinterested, disengaged, and disconnected from active play, recreational exercise, and sport activities.

Introduction

Today's children and adolescents (“youth”) are physically weaker than previous generations, and a vast majority are not accumulating at least 60 min of moderate to vigorous physical activity (MVPA) daily (1–3). Despite public health guidelines that recognize the potential physical, psychosocial, and cognitive benefits of MVPA, only one out of five adolescents meet international physical activity recommendations, which include aerobic and muscle-strengthening activities (4,5). Along with declining measures of cardiorespiratory fitness in children and adolescents (6), temporal trends indicate a steady decline in selected field measures of muscular fitness (*i.e.*, muscular strength, muscular endurance, and muscular power) in youth across the globe (7–9). Because a certain amount of force production (*e.g.*, propulsion during a jump)

The aim of this article is to discuss the interaction of neuromuscular factors that drive physical (in)activity in youth, illustrate the novel iceberg of youth physical development, and describe basic resistive skills that can be used to enhance health, improve fitness, and reduce injury risk. In this article, the term resistance training refers to an inclusive method of conditioning that involves the use of a wide range of resistance loads, different movement velocities, and varying training modalities, including body weight, free weights (barbells and dumbbells), elastic bands, weight machines, and medicine balls. As such, resistance training includes calisthenics, isometrics, circuit training, and advanced multijoint lifts.

The term dynapenia has traditionally been used to describe the loss of muscle strength that may increase the risk of physical dysfunction in older adults (19). Age-related declines in

¹Department of Kinesiology and Health Sciences, The College of New Jersey, Ewing, NJ; and ²Department of Health and Physical Education, Monmouth University, Long Branch, NJ

Address for correspondence: Avery D. Faigenbaum, EdD, FACSM, Department of Kinesiology and Health Sciences, The College of New Jersey, Ewing, NJ; E-mail: faigenba@tcnj.edu.

1537-890X/2212/414-422

Current Sports Medicine Reports

Copyright © 2023 by the American College of Sports Medicine

muscle strength and physical capacity have been linked to falls, disability, morbidity, and mortality (20). Yet the life-changing consequences of dynapenia are not limited to older adults or elderly patients with clinical conditions. Muscular weakness early in life predisposes youth to the inevitable consequences of muscle disuse, neuromuscular dysfunction, and physical inactivity (10,21,22). Even young athletes with strength deficits may be more likely to suffer a sport-related injury and less likely to reach elite level performance (23–25). Of interest, population handgrip muscle strength has been found to be an important predictor of summer Olympic games medal success in low-, middle-, and high-income countries (26).

Pediatric dynapenia can limit a child's ability to perform fundamental movement skills (FMS) proficiently and engage in exercise and sport activities with confidence, competence, and motivation (27–29). FMS include locomotor (*e.g.*, jumping), object control (*e.g.*, kicking), and stability (*e.g.*, balancing) skills that are recognized as building blocks for more advanced, complex movements (30). “Fragile” youth with strength deficits and poor movement skills may avoid nonessential physical activity to guard against failure, humiliation, and negative health outcomes, which can reduce participation in MVPA even further (16,31,32). Data from a cross-sectional study found almost 90% of children whose actual motor competence was below average did not accumulate at least 60 min of MVPA daily (33). These findings are consistent with others who reported that children with lower baseline motor performance scores maintained a stable trajectory of little to no sport participation over a 5-year study period (34).

Weaker youth increasingly engage in sedentary activities such as watching television, playing video games, or using smart phones, which are related to poor health outcomes and increased health care costs (35–37). Fraser et al. (38) compared the contributions of muscular strength measured in youth, young- and mid-adulthood with prediabetes or type 2 diabetes in midlife. They found that grip strength at each time point was equally associated with prediabetes or type 2 diabetes in mid-adulthood, with weaker participants at increased odds of poor cardiometabolic health across the life course (38). Data from a prospective cohort study of 1.2 million men found a strong association between muscular weakness during adolescence and disability 30 years later, with the strongest associations found for nervous system and psychiatric conditions (39). Others found that muscular weakness in adolescent males was inversely associated with cardiovascular disease events in middle age, independent of cardiorespiratory fitness (40).

These findings underscore the need to develop healthy, capable, and resilient youth who are able to adapt to difficult situations, withstand the demands of MVPA, and cope with life challenges (24). Because being physically strong is integral for developing and maintaining a state of resilience across the lifespan (41,42), it seems prudent to identify individuals with strength deficits early in life to reduce the burden of poor health outcomes later in life. Cut-points of muscle strength have been proposed to identify impairments in physical performance in older adults (43), and it seems similar strategies could be used to identify “at risk” youth (44). The estimated long-term economic impact of children's physical inactivity in the United States is \$1.1 trillion in direct medical costs and \$1.7 trillion in lost productivity (37). Without life course interventions that restore the loss of muscle potential, health care costs for adults

living with muscular weakness are projected to increase (45,46). Contemporary analyses of physical inactivity should recognize the critical importance of managing dynapenia in children and adults to reduce adverse health effects and the economic burden of muscle disuse (Fig. 1).

Divergence in Physical Capacity

Without early exposure to inherently enjoyable strength-building activities, current trends in MVPA will likely continue unabated as weaker youth fail to gain confidence and competence in their physical abilities (16–18). Because the decline in MVPA begins to emerge during the primary school years (about the age of 7 years) (47), interventions that prepare youth for a lifetime of physical activity should start before adolescence. Youth who do not have ongoing opportunities to improve their muscular strength may not develop the requisite physical capacities that are needed to tolerate high loads associated with exercise training and sport competition (27,48,49). This view is supported by meta-analytic data that show youth should build a foundation of strength before participating in power training activities (27).

Most children and adolescents will develop a minimal level of muscular fitness with periodic bouts of physical activity. However, regular participation in structured interventions that include resistance training (*e.g.*, physical education classes, youth fitness programs, preparatory sports conditioning) may be needed to develop robust physical capacities that optimize performance and protect against injury (50–52). Without such interventions, youth with strength deficits may be limited in their ability to generate force and explore context-specific solutions to exercise- and sport-related challenges (12,27,51). Both underweight and overweight children have been found to have impairments in muscle function, which may impact their ability to perform motor skills proficiently and engage in sport activities (53,54). An analysis of resistance-training-based sports injury prevention programs found a dose–response relationship between resistance training volume and intensity and sports injury risk reduction in participants aged 12 to 40 years (25). Of note, a relative 1 repetition maximum squat of 105% of body mass was established as the best cut-off value to distinguish high versus low risk of traumatic knee injury in adolescent female athletes (23).

Children and adolescents who enhance their muscular fitness with interventions that improve their ability to produce force may be better prepared for ongoing participation in exercise and sport activities (27,55,56). Findings from a systematic umbrella review of 14 meta-analyses found that resistance training has the potential to enhance proxies of muscular fitness in children and adolescents beyond a level achievable with growth and maturation (57). As youth enhance their muscular strength, they can build their strength reserves, which may be used (or not) to meet the workload demands of vigorous exercise and overcome unpredictable stressors on the sports field. Young athletes who continue to participate in more advanced resistance exercise programs and gradually develop their physical capacities can dramatically improve their ability to produce force, reduce force, and tolerate the workload demands of sports practice and competition (27,51,58). Conversely, those who do not participate in strength-building activities may regress along the physical capacity continuum as their muscular strength and load




Figure 1: Hypothetical representation of the costs and consequences of dynapenia throughout the life course.

tolerance begin to wane (Fig. 2). The gap between stronger and weaker youth will likely widen across developmental time without school- and community-based interventions that target strength deficits and build strength reserves (17,18). Lifestyle factors including sleep and nutrition can moderate the trajectory toward physical development or physical dysfunction (24).

Underrated Benefits of Youth Resistance Exercise

In addition to measurable improvements of muscular fitness (57,59), regular participation in youth resistance training can lead to favorable changes in musculoskeletal health, car-

diometabolic health, and mental health (60,61). Evidence supports muscular strength as a useful marker of skeletal health during development and maturation (62). High-effort resistance training (along with adequate calcium intake) can be a powerful osteogenic stimulus in youth with enduring skeletal benefits years after program completion (63,64). Higher levels of muscular fitness are associated with lower obesity and reduced cardiometabolic risk in youth (65), with evidence suggesting that low muscular fitness is causally related to the development of unfavorable levels of insulin resistance (61). Stronger youth also may be less vulnerable to mental health




Figure 2: Divergence in physical capacity among youth with varying levels of muscular strength.

disorders (42,66). Meta-analytic data indicate that resistance training can have a positive effect on some aspects of psychological well-being in youth (60).

With qualified instruction and technique-driven progression, resistance training has the potential to improve skill-related components of physical fitness, including sprinting, jumping, throwing, and agility (change of direction) (55,57,67). Because resistance-training-induced neural adaptations (*i.e.*, improvements in motor unit recruitment, firing, and coordination) can influence movement patterns in youth, regular participation in well-designed resistance training programs is likely to have a positive effect on FMS performance (57,68). These findings underscore a potential synergistic adaptation whereby training-induced gains in muscular fitness can reinforce improvements in FMS and increase participation in MVPA throughout this developmental phase of life (68–70). Emerging evidence supports a link between muscular fitness and physical activity, particularly vigorous physical activity and organized sport in children and adolescents (56).

Ongoing participation in resistance training that includes periodic bouts of higher-intensity and higher-volume lifting can enhance the load capacity of young athletes and consequently facilitate desired adaptations in musculoskeletal supporting structures (51,71,72). Stronger adult athletes are better prepared to tolerate higher workloads and larger week-to-week changes in workload than weaker athletes (73). Likewise, well-developed younger athletes with higher levels of muscular fitness may have the physical capacity to tolerate a wide range of workload demands effectively (25,71,74). Meta-analytic data from anterior cruciate ligament (ACL) injury reduction programs (which included resistance training) found an overall 50% reduction in the risk of all ACL injuries in all athletes and a 67% reduction for noncontact ACL injuries in females (75).

In light of ongoing challenges associated with adherence to injury reduction programs in youth athletes (76), early exposure to resistance training may facilitate the establishment of desired behaviors and set the stage for even greater gains in physical fitness and sport performance later in life (12,74,77). As youth learn how to perform muscle-strengthening exercises skillfully and gain confidence in their physical abilities (*i.e.*, self-efficacy), they may enhance their resistance training skill literacy as they explore different types of resistance training, learn from instructors and peers in a supporting setting, and feel good about their accomplishments (78,79). As such, resistance training skill literacy is not just about lifting weights for a prescribed number of sets and repetitions. Rather, resistance training skill literacy is an inclusive concept of knowledge, understanding, and skill performance that evolve over time to maintain interest in strength-building activities among participants with different needs, goals, and abilities (78).

All youth, especially those who are the least active, can benefit from regular participation in integrative fitness programs that include resistance training (12,80). Inactive youth are often unwilling and unable to perform prolonged periods of aerobic exercise, and participation in physical activity should not begin with competitive sport. Therefore, physical activity promotion efforts should include strength-building activities owing to its potent skill-building, performance-enhancing, and injury-reducing effects (3,57,74). An acute bout of resistance training that included body weight, medicine ball, and battling rope exercises resulted in an exercise intensity of 61% to 92%

of maximum heart rate in healthy children (81). These findings suggest that well-designed resistance training programs can enhance neuromuscular fitness and contribute to the recommended 60 min or more of MVPA daily. The underrated benefits of resistance training are becoming increasingly evident, including managing cardiometabolic risk, promoting MVPA, reducing sports-related injuries, and improving mental well-being (22,25,42,56) (Fig. 3).

Iceberg of Youth Physical Development

Years ago, Seefeldt (82) proposed that youth with poor movement skills would not be able to break through a so-called “proficiency barrier” and, therefore, would be limited in their ability to develop more mature movement patterns. Although empirical evidence supports Seefeldt’s proficiency barrier hypothesis (33,83), it appears that weaker youth with poor resistive skills may be unable to overcome a strength barrier that would allow them to perform movement skills proficiently. Because a prerequisite level of muscular strength is needed to perform a movement task skillfully (84), weaker youth may be limited in their ability (both real and perceived) to engage in MVPA with confidence, competence, and motivation (3,16). At present, global levels of FMS in children are “below average” despite the expectation that competency in these basic skills should be achieved by the age of 7 years to participate successfully in context-specific physical activities (85).

Childhood is an important period to experience the joy of moving, learn FMS, and maintain healthy developmental trajectories for future MVPA (18,34,86). The novel iceberg of physical development is a metaphoric image that graphically illustrates the sequential and cumulative influence of muscular strength on FMS and physical abilities (Fig. 4). Involuntary reflexes such as rooting and sucking and rudimentary skills that involve locomotion and manipulation provide the basis for establishing requisite levels of resistive skills as youth acquire basic movement patterns and achieve independent function. As one develops physically, more advanced movement skills that are considered foundational for ongoing participation in active play, exercise, and sport activities begin to emerge (87). The strength barrier is placed at the surface of the iceberg (the “water line”) because what is seen on the playing field is influenced by less obvious physical abilities and resistive skills developed earlier in life. As such, muscular strength can be considered a limiting factor of physical development as infants lift their body mass (*e.g.*, tummy time), toddlers grasp an external object (*e.g.*, rattle toy), preschoolers engage in outdoor activities (*e.g.*, climb a jungle gym), and children participate in active play, exercise, and sport. Importantly, ongoing participation in resistance training is needed throughout the life course to stay above the “water line” and avoid sliding down the iceberg toward a lower level of physical function.

Without adequate levels of muscular strength and neuromuscular dexterity, children may be predisposed to dynapenia and associated changes in musculoskeletal structure and function (10,21,88). Conversely, youth with adequate levels of muscular strength may be more likely to engage in vigorous physical activities, tolerate feelings and sensations associated with sport competition, and optimize gains in other important components of physical fitness (3,16,56). In support of these observations, higher levels of muscular fitness have been found to be one of the best predictors of MVPA in 5- to




Figure 3: The underrated benefits of youth resistance training.

10-year-old children (31). From a developmental perspective, stronger youth who move with confidence, competence, and motivation appear better prepared to sustain participation in MVPA across the life course (12,89). Contemporary insights into long-term physical development underscore the importance of resistance training for health improvements, performance enhancement, and injury-risk reduction (24,48,77).

Basic Resistive Skills

The importance of enhancing muscular strength to improve physical fitness was recognized by prominent leaders in early

“gymnastics-oriented” physical education (90). However, current youth guidelines that focus on the quantity of MVPA accumulated throughout the day overshadow the importance of strength-building activities (3). Efforts to enhance the physical capacity of youth should include basic resistive skills (e.g., pushing, pulling, and rotation/anti-rotation exercises) that prepare youth for more advanced forms of exercise and sport training (Fig. 5). In light of further reductions in selected measures of muscular fitness during the COVID-19 pandemic (91–93), a nuanced approach for integrating resistance training into fitness and sport programs is needed. Notably,




Figure 4: The iceberg of physical development.

misperceptions associated with youth resistance training still persist (94), and participation in muscle-strengthening activities among adolescents is falling short of expectations (95,96).

Resistance training should be integrated into school- and community-based programs to provide an important opportunity for all youth to engage in physically effortful and mentally engaging activities that enhance both health- and skill-related components of physical fitness (52,97). For example, young children could perform animal-like movements that challenge their body to function as a unit as they “frog squat,” “crocodile plank,” and “gecko lunge” (80). Child-friendly animal movements allow one to creatively paint a picture of the desired strength-building movement pattern while placing an emphasis on having fun. Resistance training with medicine balls, elastic bands, and battling ropes can be incorporated into a youth fitness circuit and individually progressed so each participant can start at a safe level and gradually progress as warranted (98). Young athletes can perform a warm-up during training sessions that includes strength-building exercises to improve neuromuscular dexterity and reduce injury risk (99). Many 5- to 7-year-olds are already involved in sport activities, and it is reasonable they also could benefit from developmentally appropriate strength-building activities (97).

The timing of brain development and its associated neuroplasticity make the early years of life an ideal time to practice basic resistive skills, learn to control posture, facilitate coordination, maintain tempo, and navigate movements efficiently (12,78). Targeting basic resistive skills that allow for

movement exploration can help to develop the requisite abilities that are needed to progress to more advanced, complex movements over time. The periodic manipulation of resistive skills as well as other program design variables (*e.g.*, sets, repetitions, rest intervals) can optimize strength gains, maintain adherence, and reduce injury risk (100). Qualified instruction, technique-driven progression, and peer support can help youth progress along the resistance training skill literacy continuum as they model desired behaviors and perform foundational movements skillfully. Detailed program design considerations are beyond the scope of this article but are available elsewhere (101).

Conclusions

Today's youth are physically weaker than previous generations and more susceptible to dynapenia and its consequences (21). Sustained reductions in muscular fitness can have significant public health consequences in terms of premature morbidity and mortality (102,103). These observations underscore the importance of enhancing muscular fitness early in life before youth possibly develop functional limitations, suffer activity-related injuries, and experience costly health conditions. If youth are not exposed regularly to interventions that include strength-building activities, they will be less likely to develop the requisite confidence and competence in their force-generating physical capabilities that are needed for ongoing participation in active play, recreational exercise, and competitive sport. A shift in our conceptual thinking about




Figure 5: Basic resistive skills.

muscular fitness is needed because established levels of physical inactivity will become more difficult to change without interventions that recognize the foundational importance of resistive skills right from the start.

The authors declare no conflict of interest and do not have any financial disclosures.

References

1. Aubert S, Barnes JD, Demchenko I, et al. Global Matrix 4.0 Physical activity report card grades for children and adolescents: results and analyses from 57 countries. *J. Phys. Act. Health.* 2022; 19:700–28.
2. Eberhardt T, Niessner C, Oriwol D, et al. Secular trends in physical fitness of children and adolescents: a review of large-scale epidemiological studies published after 2006. *Int. J. Environ. Res. Public Health.* 2020; 17:5671.
3. Faigenbaum AD, MacDonald JP, Stracciolini A, Rial Rebullido T. Making a strong case for prioritizing muscular fitness in youth physical activity guidelines. *Curr. Sports Med. Rep.* 2020; 19:530–6.
4. García-Hermoso A, López-Gil JF, Ramírez-Vélez R, et al. Adherence to aerobic and muscle-strengthening activities guidelines: a systematic review and meta-analysis of 3.3 million participants across 32 countries. *Br. J. Sports Med.* 2023; 57:225–9.
5. World Health Organization. *WHO Guidelines on Physical Activity and Sedentary Behaviour.* Geneva, Switzerland: World Health Organization; 2020. [cited 2023 Nov. 9]. Available from: <https://www.who.int/publications/i/item/9789240015128>. Accessed May 17, 2023.
6. Tomkinson GR, Lang JJ, Tremblay MS. Temporal trends in the cardiorespiratory fitness of children and adolescents representing 19 high-income and upper middle-income countries between 1981 and 2014. *Br. J. Sports Med.* 2019; 53: 478–86.
7. Kaster T, Dooley FL, Fitzgerald JS, et al. Temporal trends in the sit-ups performance of 9,939,289 children and adolescents between 1964 and 2017. *J. Sports Sci.* 2020; 38:1913–23.
8. Tomkinson GR, Kaster T, Dooley FL, et al. Temporal trends in the standing broad jump performance of 10,940,801 children and adolescents between 1960 and 2017. *Sports Med.* 2021; 51:531–48.
9. Sandercock GRH, Cohen DD. Temporal trends in muscular fitness of English 10-year-olds 1998-2014: an allometric approach. *J. Sci. Med. Sport.* 2019; 22:201–5.
10. Orsso CE, Tibaes JRB, Oliveira CLP, et al. Low muscle mass and strength in pediatrics patients: why should we care? *Clin. Nutr.* 2019; 38:2002–15.
11. Faigenbaum AD, Lloyd RS, MacDonald J, Myer GD. Citius, Altius, Fortius: beneficial effects of resistance training for young athletes: narrative review. *Br. J. Sports Med.* 2016; 50:3–7.
12. Myer G, Faigenbaum A, Edwards E, et al. Sixty minutes of what? A developing brain perspective for activating children with an integrative exercise approach. *Br. J. Sports Med.* 2015; 49:1510–6.
13. Deng Y, Fan A. Trends in sports participation in adolescents: data from a large-scale sample in the US adolescents. *Front. Public Health.* 2022; 10:960098.
14. Ryan JL, Pracht EE, Orban BL. Inpatient and emergency department costs from sports injuries among youth aged 5-18 years. *BMJ Open Sport Exerc. Med.* 2019; 5:e000491.
15. Myer GD, Jayanthi N, DiFiori JP, et al. Sports specialization, part II: alternative solutions to early sport specialization in youth athletes. *Sports Health.* 2016; 8: 65–73.
16. Zwolski C, Quatman-Yates C, Paterno MV. Resistance training in youth: laying the foundation for injury prevention and physical literacy. *Sports Health.* 2017; 9:436–43.
17. Fraser BJ, Blizzard L, Buscot M-J, et al. Muscular strength across the life course: the tracking and trajectory patterns of muscular strength between childhood and mid-adulthood in an Australian cohort. *J. Sci. Med. Sport.* 2021; 24: 696–701.
18. García-Hermoso A, Izquierdo M, Ramírez-Vélez R. Tracking of physical fitness levels from childhood and adolescence to adulthood: a systematic review and meta-analysis. *Transl. Pediatr.* 2022; 11:474–86.
19. Clark BC, Manini TM. What is dynapenia? *Nutrition.* 2012; 28:495–503.
20. Izquierdo M, Merchant RA, Morley JE, et al. International exercise recommendations in older adults (ICFSR): expert consensus guidelines. *J. Nutr. Health Aging.* 2021; 25:824–53.
21. Faigenbaum A, Rial Rebullido T, Pena J, Chulvi-Medrano I. Resistance exercise for the prevention and treatment of pediatric dynapenia. *J. Sci. Sport Exerc.* 2019; 1:208–16.
22. Gomes T, Dos Santos F, Katzmarzyk P, Maia J. Active and strong: physical activity, muscular strength, and metabolic risk in children. *Am. J. Hum. Biol.* 2017; 29.
23. Augustsson SR, Ageberg E. Weaker lower extremity muscle strength predicts traumatic knee injury in youth female but not male athletes. *BMJ Open Sport Exerc. Med.* 2017; 3:e000222.
24. Bergeron MF, Mountjoy M, Armstrong N, et al. International Olympic Committee consensus statement on youth athletic development. *Br. J. Sports Med.* 2015; 49:843–51.
25. Lauersen JB, Andersen TE, Andersen LB. Strength training as superior, dose-dependent and safe prevention of acute and overuse sports injuries: a systematic review, qualitative analysis and meta-analysis. *Br. J. Sports Med.* 2018; 52:1557–63.
26. Leong DP, McKee M, Yusuf S, PURE Investigators. Population muscle strength predicts Olympic medal tallies: evidence from 20 countries in the PURE prospective cohort study. *PLoS One.* 2017; 12:e0169821.
27. Behm DG, Young JD, Whitten JHD, et al. Effectiveness of traditional strength vs. power training on muscle strength, power and speed with youth: a systematic review and meta-analysis. *Front. Physiol.* 2017; 8:423.
28. Faigenbaum AD, MacDonald JP. Dynapenia: it's not just for grown-ups anymore. *Acta Paediatr.* 2017; 106:696–7.
29. Tveter AT, Holm I. Influence of thigh muscle strength and balance on hop length in one-legged hopping in children aged 7-12 years. *Gait Posture.* 2010; 32:259–62.
30. Logan SW, Ross SM, Chee K, et al. Fundamental motor skills: a systematic review of terminology. *J. Sports Sci.* 2018; 36:781–96.
31. Pereira S, Reyes AC, Chaves R, et al. Correlates of the physical activity decline during childhood. *Med. Sci. Sports Exerc.* 2022; 54:2129–37.
32. Sollerhed AC, Horn A, Culpán I, Lynch J. Adolescent physical activity-related injuries in school physical education and leisure-time sports. *J. Int. Med. Res.* 2020; 48:300060520954716.
33. De Meester A, Stodden D, Goodway J, et al. Identifying a motor proficiency barrier for meeting physical activity guidelines in children. *J. Sci. Med. Sport.* 2018; 21:58–62.
34. Lykkegaard CR, Andersen HS, Wehberg S, et al. The association between childhood motor performance and developmental trajectories of sport participation over 5 years in Danish students aged 6-16-year-old. *Sci. Rep.* 2023; 13:4133.
35. Carson V, Hunter S, Kuzik N, et al. Systematic review of sedentary behaviour and health indicators in school-aged children and youth: an update. *Appl. Physiol. Nutr. Metab.* 2016; 41:S240–65.
36. Hu X, Drenowatz C, Duncan M, et al. Physical education, muscle strengthening exercise, sport participation and their associations with screen time in adolescents. *Front. Public Health.* 2023; 11:1100958.
37. Lee BY, Adam A, Zenkov E, et al. Modeling the economic and health impact of increasing children's physical activity in the United States. *Health Aff. (Millwood).* 2017; 36:902–8.
38. Fraser BJ, Blizzard L, Buscot MJ, et al. The association between grip strength measured in childhood, young- and mid-adulthood and prediabetes or type 2 diabetes in mid-adulthood. *Sports Med.* 2021; 51:175–83.
39. Henriksson H, Henriksson P, Tynelius P, Ortega FB. Muscular weakness in adolescence is associated with disability 30 years later: a population-based cohort study of 1.2 million men. *Br. J. Sports Med.* 2019; 53:1221–30.
40. Timpka S, Petersson IF, Zhou C, Englund M. Muscle strength in adolescent men and risk of cardiovascular disease events and mortality in middle age: a prospective cohort study. *BMC Med.* 2014; 12:62. doi:10.1186/1741-7015-12-62.
41. Izquierdo M, Fiatarone Singh M. Promoting resilience in the face of ageing and disease: the central role of exercise and physical activity. *Ageing Res. Rev.* 2023; 88:101904.
42. Zhang X, Jiang C, Zhang X, Chi X. Muscle-strengthening exercise and positive mental health in children and adolescents: an urban survey study. *Front. Psychol.* 2022; 13:933877.
43. Tessier A-J, Wing SS, Rahme E, et al. Physical function-derived cut-points for the diagnosis of sarcopenia and dynapenia from the Canadian longitudinal study on aging. *J. Cachexia. Sarcopenia Muscle.* 2019; 10:985–99.
44. Castro-Piñero J, Perez-Bey A, Cuenca-García M, et al. Muscle fitness cut points for early assessment of cardiovascular risk in children and adolescents. *J. Pediatr.* 2019; 134:134–141.e3.
45. Pinedo-Villanueva R, Westbury LD, Syddall HE, et al. Health care costs associated with muscle weakness: a UK population-based estimate. *Calif. Tissue Int.* 2019; 104:137–44.
46. Chaput J-P, Janssen I, Sampasa-Kanyinga H, et al. Economic burden of low muscle strength in Canadian adults. *Appl. Physiol. Nutr. Metab.* 2023; 48:634–8.

47. Farooq MA, Parkinson KN, Adamson AJ, et al. Timing of the decline in physical activity in childhood and adolescence: Gateshead Millennium Cohort Study. *Br. J. Sports Med.* 2018; 52:1002–6.
48. Granacher U, Lesinski M, Busch D, et al. Effects of resistance training in youth athletes on muscular fitness and athletic performance: a conceptual model for long-term athlete development. *Front. Physiol.* 2016; 7:164.
49. Soligard T, Schweltnus M, Alonso JM, et al. How much is too much? (Part 1) International Olympic Committee consensus statement on load in sport and risk of injury. *Br. J. Sports Med.* 2016; 50:1030–41.
50. Foss KDB, Thomas S, Khoury JC, et al. A school-based neuromuscular training program and sport-related injury incidence: a prospective randomized controlled clinical trial. *J. Athl. Train.* 2018; 53:20–8.
51. Lesinski M, Prieske O, Granacher U. Effects and dose–response relationships of resistance training on physical performance in youth athletes: a systematic review and meta-analysis. *Br. J. Sports Med.* 2016; 50:781–95.
52. Villa-González E, Barranco-Ruiz Y, García-Hermoso A, Faigenbaum AD. Efficacy of school-based interventions for improving muscular fitness outcomes in children: a systematic review and meta-analysis. *Eur. J. Sport Sci.* 2023; 23:444–59.
53. Herda TJ, Ryan ED, Kohlmeier M, et al. Examination of muscle morphology and neuromuscular function in normal weight and overfat children aged 7–10 years. *Scand. J. Med. Sci. Sports.* 2018; 28:2310–21.
54. Verbecque E, Coetzee D, Smits-Engelsman B. Underweight children are agile but lack power. *BMC Pediatr.* 2022; 22:490.
55. Collins H, Booth JN, Duncan A, Fawcner S. The effect of resistance training interventions on fundamental movement skills in youth: a meta-analysis. *Sports Med. Open.* 2019; 5:17.
56. Smith JJ, Eather N, Weaver RG, et al. Behavioral correlates of muscular fitness in children and adolescents: a systematic review. *Sports Med.* 2019; 49:887–904.
57. Lesinski M, Herz M, Schmelcher A, Granacher U. Effects of resistance training on physical fitness in healthy children and adolescents: an umbrella review. *Sports Med.* 2020; 50:1901–28.
58. Suchomel TJ, Nimphius S, Stone MH. The importance of muscular strength in athletic performance. *Sports Med.* 2016; 46:1419–49.
59. Sánchez Pastor A, García-Sánchez C, Marquina Nieto M, de la Rubia A. Influence of strength training variables on neuromuscular and morphological adaptations in prepubertal children: a systematic review. *Int. J. Environ. Res. Public Health.* 2023; 20:4833.
60. Collins H, Booth JN, Duncan A, et al. The effect of resistance training interventions on ‘The Self’ in youth: a systematic review and meta-analysis. *Sports Med. Open.* 2019; 5:29.
61. García-Hermoso A, Ramírez-Campillo R, Izquierdo M. Is muscular fitness associated with future health benefits in children and adolescents? A systematic review and meta-analysis of longitudinal studies. *Sports Med.* 2019; 49:1079–94.
62. Torres-Costoso A, López-Muñoz P, Martínez-Vizcaíno V, et al. Association between muscular strength and bone health from children to young adults: a systematic review and meta-analysis. *Sports Med.* 2020; 50:1163–90.
63. Paillard T, El Hage R, Rassy NA, et al. Effects of different levels of weightlifting training on bone mineral density in a group of adolescents. *J. Clin. Densitom.* 2022; 25:497–505.
64. Thein-Nissenbaum J, Weiss DM, Kliethermes SA, Scerpella TA. Adolescent bone advantages 3 years after resistance training trial. *Pediatr. Exerc. Sci.* 2022; 35:61–9.
65. de Lima TR, Martins PC, Moreno YMF, et al. Muscular fitness and cardiometabolic variables in children and adolescents: a systematic review. *Sports Med.* 2022; 52:1555–75.
66. Cadenas-Sanchez C, Mena-Molina A, Torres-Lopez LV, et al. Healthier minds in fitter bodies: a systematic review and meta-analysis of the association between physical fitness and mental health in youth. *Sports Med.* 2021; 51:2571–605.
67. Chaabene H, Prieske O, Moran J, et al. Effects of resistance training on change-of-direction speed in youth and young physically active and athletic adults: a systematic review with meta-analysis. *Sports Med.* 2020; 50:1483–99.
68. Behringer M, Vom Heede A, Matthews M, Mester J. Effects of strength training on motor performance skills in children and adolescents: a meta-analysis. *Pediatr. Exerc. Sci.* 2011; 23:186–206.
69. Jaakkola T, Yli-Piipari S, Huotari P, et al. Fundamental movement skills and physical fitness as predictors of physical activity: a 6-year follow-up study. *Scand. J. Med. Sci. Sports.* 2016; 26:74–81.
70. Collins HM, Fawcner S, Booth JN, Duncan A. The impact of resistance training on strength and correlates of physical activity in youth. *J. Sports Sci.* 2022; 40:40–9.
71. Legerlotz K, Marzilger R, Bohm S, Arampatzis A. Physiological adaptations following resistance training in youth athletes—a narrative review. *Pediatr. Exerc. Sci.* 2016; 28:501–20.
72. Tumkur Anil Kumar N, Oliver JL, Lloyd RS, et al. The influence of growth, maturation and resistance training on muscle-tendon and neuromuscular adaptations: a narrative review. *Sports (Basel).* 2021; 9:59.
73. Malone S, Hughes B, Doran DA, et al. Can the workload-injury relationship be moderated by improved strength, speed and repeated-sprint qualities? *J. Sci. Med. Sport.* 2019; 22:29–34.
74. Petushek EJ, Sugimoto D, Stoolmiller M, et al. Evidence-based best-practice guidelines for preventing anterior cruciate ligament injuries in young female athletes: a systematic review and meta-analysis. *Am. J. Sports Med.* 2019; 47:1744–53.
75. Webster KE, Hewett TE. Meta-analysis of meta-analyses of anterior cruciate ligament injury reduction training programs. *J. Orthop. Res.* 2018; 36:2696–708.
76. Emery CA, Pasanen K. Current trends in sport injury prevention. *Best Pract. Res. Clin. Rheumatol.* 2019; 33:3–15.
77. Lloyd R, Cronin J, Faigenbaum A, et al. The National Strength and Conditioning Association position statement on long-term athletic development. *J. Strength Cond. Res.* 2016; 30:1491–509.
78. Faigenbaum A, McFarland J. Developing resistance training skill literacy in youth. *J. Phys. Ed. Rec. Dance.* 2023; 94:5–10.
79. Smith J, Diallo T, Bennie J, et al. Factors associated with adherence to the muscle-strengthening activity guideline among adolescents. *Psychol. Sport Exerc.* 2020; 51:101747.
80. Faigenbaum A, Bruno L. A fundamental approach for treating pediatric dynapenia in kids. *ACSMs Health Fit. J.* 2017; 21:18–24.
81. Faigenbaum AD, Kang J, Ratamess NA, et al. Acute cardiometabolic responses to multi-modal integrative neuromuscular training in children. *J. Funct. Morphol. Kinesiol.* 2019; 4:39.
82. Seefeldt V. Developmental motor patterns: implications for elementary school physical education. In: Nadeau K, Newell K, Roberts G, editors. *Psychol. Motor Behav. Sport.* Champaign (IL): Human Kinetics; 1980. p. 314–23.
83. Dos Santos FG, Pacheco MM, Stodden D, et al. Testing Seefeldt’s proficiency barrier: a longitudinal study. *Int. J. Environ. Res. Public Health.* 2022; 19:7184.
84. Malina R, Bouchard C, Bar-Or O. *Growth, Maturation and Physical Activity.* 2nd ed. Champaign (IL): Human Kinetics; 2004.
85. Bolger LE, Bolger LA, O’Neill C, et al. Global levels of fundamental motor skills in children: a systematic review. *J. Sports Sci.* 2021; 39:717–53.
86. O’Brien W, Khodaverdi Z, Bolger L, et al. Exploring recommendations for child and adolescent fundamental movement skills development: a narrative review. *Int. J. Environ. Res. Public Health.* 2023; 20:3278.
87. Hulteen RM, Morgan PJ, Barnett LM, et al. Development of foundational movement skills: a conceptual model for physical activity across the lifespan. *Sports Med.* 2018; 48:1533–40.
88. Behan S, Belton S, Peers C, et al. Exploring the relationships between fundamental movement skills and health related fitness components in children. *Eur. J. Sport Sci.* 2022; 22:171–81.
89. Fraser BJ, Blizzard L, Cleland V, et al. Childhood factors associated with muscular strength trajectories between childhood and mid-adulthood. *Med. Sci. Sports Exerc.* 2022; 45:1911–8.
90. Siedentop D, van der Mars H. *Introduction to Physical Education, Fitness and Sport.* 9th ed. Champaign (IL): Human Kinetics; 2023.
91. Rúa-Alonso M, Rial-Vázquez J, Nine I, et al. Comparison of physical fitness profiles obtained before and during COVID-19 pandemic in two independent large samples of children and adolescents: DAFIS project. *Int. J. Environ. Res. Public Health.* 2022; 19:3963.
92. Wahl-Alexander Z, Camic CL. Impact of COVID-19 on school-aged male and female health-related fitness markers. *Pediatr. Exerc. Sci.* 2021; 33:61–4.
93. Yuki A, Tamase Y, Nakayama M. Association between decreased grip strength in preschool children and the COVID-19 pandemic: an observational study from 2015 to 2021. *J. Physiol. Anthropol.* 2023; 42:4.
94. Faigenbaum AD, Stracciolini A, MacDonald JP, Rial Rebullido T. Mythology of youth resistance training. *Br. J. Sports Med.* 2022; bjsports-2022-105804.
95. Bennie JA, Faulkner G, Smith JJ. The epidemiology of muscle-strengthening activity among adolescents from 28 European countries. *Scand. J. Public Health.* 2022; 50:295–302.
96. Chen S, Yan J, Zhao Y. A trend analysis of adherence to the muscle strengthening exercise guidelines in US adolescents. *Int. J. Public Health.* 2022; 67:1605022.
97. Stricker PR, Faigenbaum AD, McCambridge TM, Council on Sports Medicine and Fitness. Resistance training for children and adolescents. *Pediatrics.* 2020; 145:e20201011.
98. Bukowsky M, Faigenbaum A, Myer G. FUNdamental integrative training (FIT) for physical education. *J. Phys. Ed. Rec. Dance.* 2014; 85:23–30.

99. Al Attar WS, Bizzini M, Alzahrani H, *et al.* The FIFA 11+ kids injury prevention program reduces injury rates among male children soccer players: a clustered randomized controlled trial. *Sports Health*. 2023; 15:397–409.
100. Haff G. Periodization strategies for youth development. In: Lloyd R, Oliver J, editors. *Strength and Conditioning for Young Athletes: Science and Application*. 2nd ed. Oxford (UK): Routledge Publishing; 2020. p. 281–99.
101. Faigenbaum A, Lloyd R, Oliver J, American College of Sports Medicine. *Essentials of Youth Fitness*. Champaign (IL): Human Kinetics; 2020.
102. García-Hermoso A, Cavero-Redondo I, Ramírez-Vélez R, *et al.* Muscular strength as a predictor of all-cause mortality in an apparently healthy population: a systematic review and meta-analysis of data from approximately 2 million men and women. *Arch. Phys. Med. Rehabil.* 2018; 99:2100–13.e3.
103. Momma H, Kawakami R, Honda T, Sawada SS. Muscle-strengthening activities are associated with lower risk and mortality in major non-communicable diseases: a systematic review and meta-analysis of cohort studies. *Br. J. Sports Med.* 2022; 56:755–63.